HASSE-MINKOWSKI THEOREM FOR QUADRATIC FORMS ON MORDELL-WEIL TYPE GROUPS.

STEFAN BARAŃCZUK

Abstract. In this paper we investigate an analogue of Hasse-Minkowski theorem for quadratic forms on Mordell-Weil type groups over number fields like S-units, abelian varieties with trivial ring of endomorphisms and odd algebraic K-theory groups.

arXiv:1703.06089v1 [math.NT] 17 Mar 2017

1. Introduction.

Let V denote all places on $\mathbb{Q}$ including ∞. The Hasse-Minkowski theorem (cf. [H1]) states that a quadratic form over $\mathbb{Q}$ represents 0 if and only if it represents 0 in $\mathbb{Q}_v$ for every $v \in V$. Since quadratic forms can be defined on any module over a commutative ring it is a natural question whether we can obtain an analogue of Hasse-Minkowski theorem provided our module is equipped with a system of reduction maps.

A perfect example of such module is the Mordell-Weil group of an elliptic curve over a number field. Observe that here not every place defines a reduction map since we have to exclude primes of bad reduction. Note however that the original Hasse-Minkowski theorem also has its stronger versions where some number of places can be omitted (see Proposition 4); our proofs depend on those results.

The groups for which we prove Hasse-Minkowski theorem in this paper are the following Mordell-Weil type groups:

(1) $R_{F,S}^{\times}$, S-units groups, where F is a number field and S is a finite set of ideals in the ring of integers R_F,
(2) $A(F)$, Mordell-Weil groups of abelian varieties over number fields F with $\mathrm{End}_{\bar{F}}(A) = \mathbb{Z}$,
(3) $K_{2n+1}(F)$, $n > 0$, odd algebraic K-theory groups.

To deal with them it is enough for us to introduce the following abstract nonsense axiomatic setup:

Let B be an abelian group and $r_v \colon B \to B_v$ be an infinite family of groups homomorphisms whose targets B_v are finite abelian groups. We will use the following notation:

$P \mod v$	denotes $r_v(P)$ for $P \in B$
$P = Q \mod v$	means $r_v(P) = r_v(Q)$ for $P, Q \in B$
Λ_{tors}	the torsion part of a subgroup $\Lambda < B$
$\mathrm{ord}\, T$	the order of a torsion point $T \in B$
$\mathrm{ord}_v P$	the order of a point $P \mod v$
$l^k \parallel n$	means that l^k exactly divides n, i.e. $l^k \mid n$ and $l^{k+1} \nmid n$ where l is a prime number, k a positive integer and n a natural number.

We impose the following two assumptions on the family $r_v \colon B \to B_v$:

Assumptions.

(1) Let l be a prime number and $(k_1, \ldots, k_m)$ a sequence of nonnegative integers. If $P_1, \ldots, P_m \in B$ are points linearly independent over $\mathbb{Z}$ then there is a positive

2010 *Mathematics Subject Classification.* 11R04; 11R70; 14K15.
Key words and phrases. quadratic forms; Hasse-Minkowski theorem; Mordell-Weil groups.

density set of primes v in F such that $l^{k_i} \parallel \mathrm{ord}_v P_i$ if $k_i > 0$ and $l \nmid \mathrm{ord}_v P_i$ if $k_i = 0$.

(2) For almost all v the map $B_{\mathrm{tors}} \to B_v$ is injective.

(For the families of groups we described above the axioms are valid, in particular Assumption 1 is fulfilled by [Bar], Theorem 5.1 and Assumption 2 by [BGK], Lemma 3.11. The validity of the technical hypothesis [1] from the first sentence of Theorem 5.1 in [Bar] is clear in the K-theory groups case (resp. $R^{\times}_{F,S}$ case) since here ρ_l is simply one-dimensional representation given by the $(n+1)$th tensor power of cyclotomic character (resp. by cyclotomic character); in abelian varieties case it is asserted by Corollary 1 in [Bog].)

2. Quadrartic forms of rank 2.

Theorem 1. *Let $P, Q \in B$ be points of infinite order. The following are equivalent:*

- *For almost every v there exist coprime integers x, y and a point $T \in B_{\mathrm{tors}}$ such that*

$$x^2 P + y^2 Q = T \mod v. \tag{1}$$

- *There exist coprime integers x, y and a point $T \in B_{\mathrm{tors}}$ such that*

$$x^2 P + y^2 Q = T. \tag{2}$$

Proof. Suppose that P, Q are linearly independent. Fix a prime number l coprime to $\#B_{\mathrm{tors}}$. By Assumption 1 there is a positive density set of primes v such that

$$l^2 \parallel \mathrm{ord}_v P \quad \text{and} \quad l^3 \parallel \mathrm{ord}_v Q.$$

Thus if both x, y are coprime to l then

$$l^2 \parallel \mathrm{ord}_v(x^2 P) \quad \text{and} \quad l^3 \parallel \mathrm{ord}_v(y^2 Q);$$

if $l \mid x$ and $l \nmid y$ then

$$l \nmid \mathrm{ord}_v(x^2 P) \quad \text{and} \quad l^3 \parallel \mathrm{ord}_v(y^2 Q);$$

and if $l \nmid x$ and $l \mid y$ then

$$l^2 \parallel \mathrm{ord}_v(x^2 P) \quad \text{and} \quad l^2 \nmid \mathrm{ord}_v(y^2 Q);$$

in all cases we get a contradiction to (1). Hence P, Q are linearly dependent.

Write $aP + bQ = 0$ for nonzero integers a, b. Multiplying (1) by b we get

$$(bx^2 - ay^2)P = bT \mod v. \tag{3}$$

Fix an arbitrary prime number l coprime to $\#B_{\mathrm{tors}}$. By Assumption 1 there is a positive density set of primes v such that $l \parallel \mathrm{ord}_v P$. So by (3) there are coprime integers x, y such that $l \mid (bx^2 - ay^2)$. Thus by Proposition 4 (a) the form $bx^2 - ay^2$ represents 0. Putting $bx^2 - ay^2 = 0$ we get $b(x^2 P + y^2 Q) = 0$.

□

3. Quadrartic forms of rank 3.

Theorem 2. *Let $P, Q, R \in B$ be points of infinite order. The following are equivalent:*

- *For almost every v there exist integers x, y, z with $\gcd(x, y, z) = 1$ and a point $T \in B_{\mathrm{tors}}$ such that*

$$x^2 P + y^2 Q + z^2 R = T \mod v. \tag{4}$$

- *There exist integers x, y, z with $\gcd(x, y, z) = 1$ and a point $T \in B_{\mathrm{tors}}$ such that*

$$x^2 P + y^2 Q + z^2 R = T. \tag{5}$$

[1] note the misprint in its formulation: $\rho(G_F)$ should read $\rho_l(G_F)$

Proof of Theorem 2. First we prove that P, Q, R are linearly dependent. Indeed, assume that P, Q, R are linearly independent. This means that the points $P+Q, Q, Q+R$ are also independent. Let 2^e be the maximal power of 2 dividing $\#B_{\mathrm{tors}}$. By Assumption 1 there is a positive density set of primes v such that

$$2^{1+e} \parallel \mathrm{ord}_v(P+Q) \quad , \quad 2^{3+e} \parallel \mathrm{ord}_v Q \quad , \quad 2^{4+e} \parallel \mathrm{ord}_v(Q+R). \tag{6}$$

That gives

$$2^{3+e} \parallel \mathrm{ord}_v P \quad , \quad 2^{3+e} \parallel \mathrm{ord}_v Q \quad , \quad 2^{4+e} \parallel \mathrm{ord}_v R. \tag{7}$$

Now by (4) we have $2 \mid z$ thus

$$\text{either} \quad 2^{1+e} \nmid \mathrm{ord}_v(z^2 R) \quad \text{or} \quad 2^{2+e} \parallel \mathrm{ord}_v(z^2 R) \tag{8}$$

so respectively

$$\text{either} \quad 2^{1+e} \nmid \mathrm{ord}_v(x^2 P + y^2 Q) \quad \text{or} \quad 2^{2+e} \parallel \mathrm{ord}_v(x^2 P + y^2 Q). \tag{9}$$

Since $\gcd(x, y, z) = 1$ we have to consider two cases: when both x, y are odd and when exactly one of x, y is even.

If both x, y are odd then $8 \mid (x^2 - y^2)$ so we get by (6) and (7) that

$$2^{1+e} \parallel \mathrm{ord}_v(y^2(P+Q)) \quad \text{and} \quad 2^{1+e} \nmid \mathrm{ord}_v((x^2-y^2)P).$$

Since $x^2 P + y^2 Q = y^2(P+Q) + (x^2 - y^2)P$ we have $2^{1+e} \parallel \mathrm{ord}_v(x^2 P + y^2 Q)$ but that contradicts (9).

If x is odd and y is even then

$$2^{3+e} \parallel \mathrm{ord}_v(x^2 P) \quad \text{and} \quad 2^{3+e} \nmid \mathrm{ord}_v(y^2 Q)$$

but that together with (8) contradicts (4). If y is odd and x is even then we argue in the same manner.

We have proven that P, Q, R are linearly dependent.

First we analyse the case when among the pairs $\{P, Q\}$, $\{P, R\}$, $\{Q, R\}$ one is a pair of linearly dependent points. Without loss of generality we can assume that this pair is $\{P, Q\}$, i.e., there are nonzero rational integers s, t such that $sP = tQ$. If $-s/t$ is the square of a rational number then we are done. Indeed, write $-s/t = (p/q)^2$ with coprime p, q. Then $t(p^2 P + q^2 Q) = 0$ so we can put $(x, y, z) = (p, q, 0)$ in (5).

So suppose that $-s/t$ is not a square. Multiplying (4) by s we get

$$(tx^2 + sy^2)Q + sz^2 R = sT \mod v. \tag{10}$$

Fix a prime number l such that l is coprime to both s, t and to $\#B_{\mathrm{tors}}$ and such that $-s/t$ is a quadratic nonresidue modulo l. Suppose that Q, R are linearly independent. By Assumption 1 there is a positive density set of primes v such that $l^2 \parallel \mathrm{ord}_v Q$ and $l^3 \parallel \mathrm{ord}_v R$. Then (10) implies that $l \mid z$. This means that $l^2 \nmid \mathrm{ord}_v(sz^2 R)$ so again by (10) we get $l \mid (tx^2 + sy^2)$. But $\gcd(x, y, z) = 1$ hence both x, y are coprime to l and we can write $(x/y)^2 = -s/t \mod l$ contrary to our assumption that $-s/t$ is nonresidue. Thus Q, R are linearly dependent.

Write $uQ = wR$ for some nonzero rational integers u, w. Multiplying (10) by w we get

$$(wtx^2 + wsy^2 + usz^2)Q = wsT \mod v. \tag{11}$$

Denote by $\mathcal{P}$ the set of all prime divisors of $\#B_{\mathrm{tors}}$. For every $p \in \mathcal{P}$ let p^{e_p} be the maximal power of p dividing $\#B_{\mathrm{tors}}$. Define $e = \max\{e_p : p \in \mathcal{P}\}$. Fix an arbitrary prime number l and a positive integer k. By Assumption 1 there is a positive density set of primes v such that $l^{k+e} \parallel \mathrm{ord}_v Q$ so $l^k \mid (wtx^2 + wsy^2 + usz^2)$ by (11). Thus the

quadratic form $(wtx^2 + wsy^2 + usz^2)$ represents 0 modulo arbitrary l^k and hence by Proposition 4 (b) it represents 0.

Now we turn to the case when among the pairs $\{P, Q\}$, $\{P, R\}$, $\{Q, R\}$ none is a pair of linearly dependent points.

Write

$$aP + bQ + cR = 0 \tag{12}$$

with $abc \neq 0$. Multiplying (4) by a and (12) by x^2 and subtracting the results we get

$$(ay^2 - bx^2)Q + (az^2 - cx^2)R = aT \mod v. \tag{13}$$

Fix a prime number l coprime to the numbers $a, b, c, \#B_{\text{tors}}$. By Assumption 1 there is a positive density set of primes v such that

$$l^2 \parallel \operatorname{ord}_v Q \quad \text{and} \quad l \parallel \operatorname{ord}_v R. \tag{14}$$

By (13) we get

$$l \mid (ay^2 - bx^2) \tag{15}$$

thus if l divides either x or y then it divides them both. But that implies by (14) and (13) that $l \nmid \operatorname{ord}_v(az^2R)$ so we have $l \mid z$ which is a contradiction to our assumption that $\gcd(x, y, z) = 1$. Thus l divides neither x nor y and (15) means that b/a is a square modulo l.

The only restriction on l in the above argument is that it cannot divide the numbers $a, b, c, \#B_{\text{tors}}$ hence it is valid for all but finitely many prime numbers l and we can use Proposition 4 (a) which asserts that b/a is the square of a rational number. Analogously we prove the same for c/a.

Hence we can put $b/a = (t/s)^2$ and $c/a = (u/w)^2$ and by (12) we get

$$a(sw)^2P + a(tw)^2Q + a(su)^2R = 0.$$

□

4. Quadratic forms of higher ranks. Auxiliary results.

Theorems 1 and 2 cannot be extended directly for quadratic forms of higher ranks, even if we assume that they represent zero for all primes of good reduction. Indeed, we have the following result:

Proposition 3. *Fix $n \geq 4$. Let P be a point of infinite order. Define the form f by*

$$(2x_1^2 + x_2^2 + x_3^2 + x_4^2 + \ldots + x_n^2)P.$$

Then $f = 0$ if and only if $(x_1, \ldots, x_n) = (0, \ldots, 0)$ but for every prime v of good reduction there exist integers $x_1, \ldots, x_n$ with $\gcd(x_1, \ldots, x_n) = 1$ such that

$$(2x_1^2 + x_2^2 + x_3^2 + x_4^2 + \ldots + x_n^2)P = 0 \mod v. \tag{16}$$

Proof. The form $(2x_1^2 + x_2^2 + x_3^2 + x_4^2 + \ldots + x_n^2)$ is positive and the point P is of infinite order so $(2x_1^2+x_2^2+x_3^2+x_4^2+\ldots+x_n^2)P = 0$ if and only if $(x_1, \ldots, x_n) = (0, \ldots, 0)$.

Let k be the order of $P \mod v$. It is an easy corollary of the theorem of Gauss on sums of three squares (cf. [Sie], page 395, exercise 4) that $2k$ is of the form $2a^2+b^2+c^2+1$ where a, b, c are integers. Thus in order to get (16) we can put $x_1 = a, x_2 = b, x_3 = c, x_4 = 1$ and $x_i = 0$ for $i > 4$. □

Proposition 4.

(a) *Let f be an integral quadratic form of rank 2. If $f \mod v$ represents 0 for all but finitely many $v \in V$ then f represents 0.*

(b) *Let f be an integral quadratic form of rank* 3. *If f represents* 0 *in $\mathbb{Q}_v$ for all except at most one $v \in V$ then f represents* 0.

Proof. (a) This statement is an immediate corollary of the theorem proved in [H2].
(b) Cf. [S], p. 43, Corollary 3. □

References

[BGK] G. Banaszak, W. Gajda, P. Krasoń, *Detecting linear dependence by reduction maps*, Journal of Number Theory 115 (2005), 322-342

[Bar] S. Barańczuk, *On reduction maps and support problem in K-theory and abelian varieties*, Journal of Number Theory 119 (2006), 1-17

[Bog] F.A. Bogomolov, *Sur l'algébricité des représentations l-adiques*, C. R. Acad. Sci. Paris Sr. A-B 290 (1980), no. 15, A701-A703

[H1] H. Hasse, *Über die Darstellbarkeit von Zahlen durch quadratische Formen im Körper der rationalen Zahlen*, J. Reine Angew. Math. 152 (1923), 129-148

[H2] H. Hasse, *Zwei Bemerkungen zu der Arbeit "Zur Arithmetik der Polynome" von U. Wegner in den Mathematischen Annalen, Bd. 105, S. 628-631*, Math. Ann. 106 (1932), no. 1, 455-456

[S] J. -P. Serre, *A course in Arithmetic*, Graduate Texts in Mathematics, Springer 1996

[Sie] W. Sierpiński, *Elementary Theory of Numbers*, North-Holland mathematical library, vol. 31, 1988

Faculty of Mathematics and Computer Science, Adam Mickiewicz University, ul. Umultowska 87, Poznań, Poland

E-mail address: stefbar@amu.edu.pl